

EXERCISE 2

Backward “Hello World” NPP example, Forward NPP example

Aims:

1. Backward:


1. Familiarise with FLEXPART run in backward mode
2. Understand the strengths and limits of backtracking a single measurement – poor man’s inversion

2. Forward:

1. How to set a realistic multi-species scenario
2. Influence of grid, depositing characteristics
3. Thinking towards emergency response

Questions? Write delia.arnold-arias@zamg.ac.at

Sites of Nuclear Power Plants in China



- 1 day integration measurement of Xe-133 (can be treated as a tracer) at the HKO
22.3025° N, 114.1742° E
- Measurement integration period 2014.10.12 00:00 - 2014.10.13 00:00
- Measurement inlet at 60 m a.g.l.
- Backtracking for 3 days
- Interested in potential emissions at ground level, therefore OUTGRID will only need one level, relatively shallow, 150 m a.g.l.
- Remember to carefully check the COMMAND
- Output every three hours

2014.10.12 00:00 - 2014.10.13 00:00 – Measurement at
HKO! $4 \times 10^{**4}$ Bq/m³ Xe-133

This test case aims at making you run FLEXPART with a simple bwd case and understand the results and what information can be extracted from them.

■ Specifications:

■ COMMAND FILE:

- Backward run: 9 October 2014 00 UTC to 13 October 2014 00 UTC
- Output every 3 hours
- Convection
- Residence times output
- No nested output
- No adaptation to TL

■ OUTGRID FILE:

- Resolution 0.2 degree
- LLC: 20.0° N, 100.0° E
- 100x150 grid cells
- 1 output layer, 150 m agl

■ RELEASES FILE:

- Point source at the top of the HKO building: 22.3025° N, 114.1742° E, 60 m agl
- Release start: 12 October 2014 00 UTC
- Release end: 13 October 2014 00 UTC
- Total mass: 1 kg of Xe-133
- Particles released: 10000

- 1 day integration measurement of Xe-133 (can be treated as a tracer) at the HKO
22.3025° N, 114.1742° E
- Measurement integration period 2014.10.12 00:00 - 2014.10.13 00:00
- Measurement inlet at 60 m a.g.l.
- Backtracking for 3 days before measurement start
- Interested in potential emissions at ground level, therefore OUTGRID will only need one level, relatively shallow, 150 m a.g.l.
- Remember to carefully check the COMMAND
- Output every three hours

```
Welcome to FLEXPART Version 10.3beta (2019-04-05)
FLEXPART is free software released under the GNU General Public License.
-----
INFORMATION: SUBGRIDSCALE TERRAIN EFFECT IS
NOT PARAMETERIZED DURING THIS SIMULATION.
-----
ECMWF metdata detected
Vertical levels in ECMWF data: 138 138

Mother domain:
Longitude range: 100.00000 to 130.00000 Grid distance: 0.20000
Latitude range : 20.00000 to 40.00000 Grid distance: 0.20000

Releasepoints : 1


SPECIES: 21 Xe-133 (GAS)
Wet removal for gases is turned: OFF
Dry deposition for gases is turned: OFF
Below-cloud scavenging: OFF
In-cloud scavenging: OFF
Particles allocated (maxpart) : 5000000
Particles released (numpartmax): 10000
Total mass released: 1.0000000E+00
Allocating fields for global output (x,y): 150 100
Allocating fields for nested output (x,y): 0 0
Concentrations are calculated using kernel
Simulated 0.0 hours ( 0 s), 0 particles
Global ECMWF fields: using cloud water
Global ECMWF fields: using cloud water
Global ECMWF fields: using cloud water
-10800 Seconds simulated: 1302 Particles: Uncertainty: 0.000 0.000 0.000
Global ECMWF fields: using cloud water
-21600 Seconds simulated: 2552 Particles: Uncertainty: 0.000 0.000 0.000
```

ECMWF (0.2 degr.) : Bwd run from HKO

```
./plot_FLEX_binary.py ./output/ False nuc1 0 0 alldates cyl False
```


```
100.0,130.0,20.0,40.0 mesh False
```

[In backward mode all species are treated the same: write nuc1 or tracer1.]



NCEP (0.5 degr.): Bwd run from HKO




```
./plot_FLEX_binary.py ./output_NCEP/ False nuc1 0 0 alldates cyl False  
100.0,130.0,20.0,40.0 mesh False
```



Is this discontinuous plume evolution reasonable? Remember differences between COMPUTATIONAL domain and OUTGRID domain

Bwd run from HKO




Sites of Nuclear Power Plants in China



Do we have only one single potential contributing source?

Bwd run from HKO

Sites of Nuclear Power Plants in China



What would be the sources that would have led to that measurement with respect to the results?

Observation = Emission * sensitivity to emission
 $m[Bq/m^3] = M[s] * S[Bq/s * m^3]$

How many Bequerels (kg, ...) released?

Potential source 1 – Daya Bay

M?

How big is the grid size?



Which is the average timing? In what units?

Potential source 2 – Qinshan

M?

How big is the grid size?

Which is the average timing? In what units?



What would be the sources that would have led to that measurement with respect to the results?

Observation = Emission * sensitivity to emission
 $m[Bq/m^3] = M[s] * S[Bq/s \cdot m^3]$

How many Bequerels (kg, ...) released?

Potential source 1 – Daya Bay

M? 0.15 s

How big is the grid size? 0.2 x 0.2 deg x 150 (m)



Which is the average timing? In what units? 3 hours

Potential source 2 – Qinshan

M?

How big is the grid size? 0.2 x 0.2 deg x 150 (m)

Which is the average timing? In what units? 3 hours



What would be the sources that would have led to that measurement with respect to the results?

Observation = Emission * sensitivity to emission
 $m[\text{Bq}/\text{m}^3] = M[\text{s}] * S[\text{Bq}/\text{s} * \text{m}^3]$

How many Bequerels (kg, ...) released?

Potential source 1 – Daya Bay

M? 0.15 s

How big is the grid size? 0.2 x 0.2 deg x 150 (m)

Which is the average timing? In what units? 3 hours

$$S[\text{Bq}] = (m[\text{Bq}/\text{m}^3]/M[\text{s}]) * 3 * 3600[\text{s}] * 0.2 * 100000 * 0.2 * 100000 * 150[\text{m}^3]$$

$$= 4 * 10^{14} [\text{Bq}/\text{m}^3] * 1/M[\text{s}] * 3 * 3600[\text{s}] * 0.2 * 100000$$

$$* 0.2 * 100000 * 150[\text{m}^3] = 1.77\text{E}+20 \text{ Bq!}$$

Potential source 2 – Qinshan

M? 3.5 s



How big is the grid size? 0.2 x 0.2 deg x 150 (m)

Which is the average timing? In what units? 3 hours

$$S[\text{Bq}] = (m[\text{Bq}/\text{m}^3]/M[\text{s}]) * 3 * 3600[\text{s}] * 0.2 * 100000 * 0.2 * 100000 * 150[\text{m}^3]$$


$$= 4 * 10^{14} [\text{Bq}/\text{m}^3] * 1/M[\text{s}] * 3 * 3600[\text{s}] * 0.2 * 100000$$

$$* 0.2 * 100000 * 150[\text{m}^3] = 7.41\text{E}+18 \text{ Bq!}$$



?

Sites of Nuclear Power Plants in China



2 reactor units
 Framatome 900 MWe
 PWR

GNPS NPP hypothetical scenario

Table 3.4-1 Hypothetical source term used in the contingency plan in Hong Kong

Source term GD	Probability (y ⁻¹)	Release time (h)	Release duration (h)	Height (m)	Release fraction						
					Xe,Kr	I	Cs	Te	Ba,Sr	Ru ⁽¹⁾	La ⁽²⁾
1	3.00E-08	1.5	1	10	2.0	1.6	1.44	1.38	1.1E-1	6.6E-1	1.7E-2
2	5.60E-06	1.5	1	10	1.0	8.0E-1	7.2E-1	6.9E-1	5.6E-2	3.3E-1	8.7E-3
3	2.00E-07	2.5	1	10	1.0	1.8E-1	1.4E-1	1.4E-1	5.0E-2	5.8E-3	6.4E-4
4	2.60E-07	2.5	1	10	1.0	1.8E-1	1.4E-1	1.4E-1	5.0E-2	5.8E-3	6.4E-4
5	1.20E-06	1	10	10	1.0	1.8E-1	8.4E-2	3.6E-1	1.3E-1	2.0E-2	1.8E-3
6	4.00E-07	1	10	10	1.0	5.2E-2	1.7E-2	7.2E-2	2.6E-2	4.1E-3	3.6E-4
7	4.80E-05	12	1	10	1.0	1.7E-1	1.5E-1	2.4E-1	5.3E-2	4.2E-5	3.4E-3
8	6.00E-08	2.5	10	10	1.0	5.0E-2	3.0E-2	3.7E-2	2.1E-2	2.2E-3	8.9E-3
9	1.20E-06	24	1	0	1.0	1.7E-4	1.0E-4	9.7E-5	1.1E-5	8.9E-7	1.3E-7
10	1.20E-06	24	10	10	0.75	8.5E-3	3.5E-3	3.5E-3	4.0E-3	3.0E-4	5.0E-5
11	5.00E-06	1	2	0	1.0	6.7E-1	6.7E-1	6.4E-1	7.0E-2	5.0E-2	9.0E-3
12	3.60E-05	24	1	10	5E-3	1.2E-4	1.0E-4	9.7E-5	1.1E-5	8.9E-7	1.3E-7

(1) include Ru, Mo (2) include La, Y, Zr, Ce, Pu, Am, Cm

Source from:

Consultancy on the environment aspects of the Daya Bay Nuclear Power Station for the government of Hong Kong – Risk Assessment [16]

<http://hub.hku.hk/bitstream/10722/41313/6/FullText.pdf>

Hypothetical ST based on US Surry (Westinghouse) PWR due to similar design (not the same thermal power – scaling)




Table 3.4-2 Description of the source term used in the hypothetical accident

GD	Description
1, 2	Early overpressure due to direct heating, hydrogen explosion, or steam explosion effects, or combinations thereof, with high pressure
3	Early overpressure due to direct heating, hydrogen explosion, or steam explosion effects, or combinations thereof, with low or medium pressure core melt and sprays NOT operating.
4	Early overpressure due to direct heating, hydrogen explosion, or steam explosion effects, or combinations thereof, with low or medium pressure core melt and sprays operating.
5	Failure to isolate (large leakage rate), sprays NOT operating.
6	Failure to isolate (large leakage rate), sprays operating.
7	Overpressure due to failure of containment heat removal.
8	Delayed overpressure due to accumulation of incondensable gases, or delayed excessive steam generation (with containment heat removal operating)
9	Basemat penetration by core debris.
10	Deliberate (filtered) venting.
11	Bypass of the containment by direct connection of the primary circuit to the surroundings
12	Successful containment.


Table 3.5-1 Inventory in reactor of the nuclear power plant and half-lives

Nuclide	Initial inventory Q_T (Bq)	Half-lives $T_{1/2}$ (s)
Kr-85	1.8490E+16	3.3860E+08
Kr-85m	7.9000E+17	1.6128E+04
Kr-87	1.4520E+18	4.5602E+03
Kr-88	1.9950E+18	1.0083E+04
Sr-89	2.8760E+18	4.4928E+06
Sr-90	1.4910E+17	8.8646E+08
Y-91	3.8550E+18	5.0553E+06
Zr-95	5.0260E+18	5.6592E+06
Mo-99	5.5000E+18	2.3760E+05
Ru-103	4.4810E+18	3.4206E+06
Ru-106	2.7210E+18	3.1795E+07
Te-129m	1.4230E+17	2.9030E+06
Te-131m	2.9620E+17	1.0800E+05
Te-132	4.1440E+18	2.8080E+05
I-131	2.8950E+18	6.9466E+05
I-132	4.2410E+18	8.2800E+03
I-133	5.8220E+18	7.4822E+04
I-134	6.6910E+18	3.1560E+03
I-135	5.4740E+18	2.3796E+04
Xe-133	5.8480E+18	4.5706E+05
Xe-135	1.4230E+18	3.3013E+04
Cs-134	3.2920E+17	6.5007E+07
Cs-136	1.1330E+17	1.1318E+06
Cs-137	1.9560E+17	9.4954E+08
Ra-140	5.2480E+18	1.1051E+06

RELEASES

RN	Inventory (Bq)	Rel. Frac.	Release (Bq)	Release (TBq)
Xe-133 NG	5.85E+18	1	5.85E+18	5.85E+06
Cs-137 aerosol	1.96E+17	0.14	2.74E+16	2.74E+04
I-131 (I2)	2.90E+18	0.18	2.61E+17	2.61E+05
I-131 aerosol			2.61E+17	2.61E+05

- Emission/release length : 1 hour starting at 20140923 00:00 UTC
- Emission/release height : 10 m a.g.l → 50 m a.g.l column (~ building height)
- Location: 22.5972° N , 114.5444° E




Multispecies release

Mind:

- Gravitational settling
- Definition of each of the species
- Post-processing
- Number of particles per release (10000)

OUTGRID/OUTGRID_NEST

- Mother/Nest:
 - LLC -179.0° / 112.5° E, -90.0° / 21.0° N
 - 360 x 180 / 320 x 320
- Vertical levels
 - 1000, 2000 and 5000 m a.g.l
- Horizontal resolution mother/nest: 1.0° x 1.0° / 0.01° x 0.01°



COMMAND

- Simulation start / stop: 20140923 00:00 UTC / 20140923 12:00 UTC
- Temporal resolution: hourly
- LCONVECTION=1, IOUT=0, NESTED_OUTPUT=1


ECMWF (1.0° / 0.2° nested) & GFS (0.5°) meteorological data

GNPS NPP hypothetical scenario


High-resolution nest needed in the near-range!

Results for
Xe-133
(non-
depositing)


ECMWF:



NCEP:



```
./plot_FLEX_binary.py
./output_ECMWF [NCEP]/ True
[False] 4 0 3 alldates cyl False
112.5,115.5,21.0,24.0 mesh
[contour] False
```



GNPS NPP hypothetical scenario


Cs-137 aerosol:

Concentration

Wet depo

Dry depo

Total depo



```
./plot_FLEX_binary.py ./output_ECMWF[NCEP]/ True 3 0 0 alldates cyl True  
112.5,115.5,21.0,24.0 contour False
```